

Double exchange model in cubic vanadates

Krzysztof Wohlfeld and Andrzej M. Oleś

Marian Smoluchowski Institute of Physics, Jagellonian University,
Reymonta 4, PL-30059 Kraków, Poland

Abstract

We investigate the role of the double exchange mechanism for stability of the metallic *C*-type antiferromagnetic (*C*-AF) phase, which was observed experimentally in hole doped $\text{La}_{1-x}\text{Sr}_x\text{VO}_3$. The double exchange model treats localized electrons in *xy* orbitals as classical $S = 1/2$ spins, which interact by Hund's exchange J_H with *yz/zx* electrons in partly filled t_{2g} orbitals. Including strong on-site Coulomb repulsion U between t_{2g} electrons, and using slave boson method we demonstrate that *C*-AF and metallic phase can be stabilized due to the specific features of the hopping in degenerate and partly filled t_{2g} orbitals.

1 Introduction

Recently undoped vanadium oxides, such as LaVO_3 or YVO_3 , have attracted much attention due to their puzzling magnetic properties [1]. Large on-site Coulomb repulsion $U \simeq 5$ eV between $3d$ electrons leads to suppression of charge fluctuations and to the effective superexchange (SE) interaction $\propto J = 4t^2/U$ in the low energy range, where $t \simeq 0.2$ eV is the hopping element. The cubic vanadates belong to the class of Mott insulators with orbital degeneracy as two d electrons, localized on the V^{3+} ion, occupy two out of three (almost) degenerate t_{2g} orbitals. Thus, the SE model includes t_{2g} orbital degrees of freedom [2], and *by itself* explains the observed *C*-type antiferromagnetic (AF) phase in LaVO_3 , with ferromagnetic (FM) chains staggered in two other directions.

Even more interesting and not yet understood are the properties of doped $\text{La}_{1-x}\text{Sr}_x\text{VO}_3$, where the following phases were experimentally observed: *C*-AF and insulating one up to $x = 0.178$ hole concentration, *C*-AF metallic one for $0.178 < x < 0.26$, and, finally, paramagnetic and metallic one for $0.26 < x < 0.327$ [3]. Theoretical explanation of the existence of this variety of phases has to take into account that the motion of doped holes is strongly affected by: (*i*) intersite SE interaction $\propto J$, (*ii*) on-site Hund's interaction $\propto J_H = 0.68$ eV [4]. Such a theoretical model was recently discussed by Ishihara [5], but there the Hund's term was introduced in a nonsystematic way via certain projection operators, with $J_H \rightarrow \infty$ limit being assumed implicitly, and the model was solved in a way suitable for small doping only. Here we start with an appropriate Kondo-lattice model, systematically

introduce the large J_H limit in which the kinetic and the Hund's parts of the Hamiltonian reduce to the double exchange (DE) model [6], and solve it for any realistic value of doping. Since in this paper we are mainly concerned with the explanation of the existence of the metallic phase and the associated magnetic order, we will simplify the SE part of the Hamiltonian as much as possible and investigate only the competition of the DE and the generic SE interaction due to the occupied xy orbitals ($n_{xy} = 1$), leaving the study of the full Hamiltonian as a future problem. In particular we want to answer three questions: (i) can C -AF and metallic phase be predicted by DE, (ii) if yes, then why such a phase contradicts the "intuitive" picture of the DE mechanism which supports homogenous FM phase, and (iii) in what way the generic features of degenerate t_{2g} orbitals help to understand the answers obtained in points (i) and (ii), especially the differences to the case of the e_g electrons, playing a role in the manganites [7]. Thereby, we shall use a similar approximation to that employed in manganites [8], assuming localized xy electrons and treating them as "core spins".

2 Double exchange Hamiltonian and the stability of the C -AF order

For the sake of clarity we start with a FM Kondo-lattice Hamiltonian but we already include important simplifications in the SE part:

$$\begin{aligned} \mathcal{H} = & \mathcal{P} \left\{ -t \sum_{i,j \parallel \hat{y}, \hat{z}} a_{i\sigma}^\dagger a_{j\sigma} - t \sum_{i,j \parallel \hat{x}, \hat{z}} b_{i\sigma}^\dagger b_{j\sigma} - J_H \sum_{\substack{i \\ \sigma, \sigma'}} \mathbf{S}_i \cdot \vec{\sigma}_{\sigma\sigma'} (a_{i\sigma}^\dagger a_{i\sigma'} + b_{i\sigma}^\dagger b_{i\sigma'}) \right\} \mathcal{P} \\ & + J \sum_{\langle ij \rangle \parallel \hat{x}, \hat{y}} \mathbf{S}_i \cdot \mathbf{S}_j, \end{aligned} \quad (1)$$

where: $\mathbf{S}_i$ are "core spin" $S = 1/2$ operators of t_{2g} electrons in occupied xy orbitals, and $\vec{\sigma}_{\sigma\sigma'}$ is a vector of Pauli matrices. The fermion operators $\{a_{i\sigma}^\dagger, b_{i\sigma}^\dagger\}$ create an electron with spin σ in yz and zx orbitals, labeled here as a and b as these orbitals have no amplitude along a and b axis [2], respectively. In the relevant regime of large Coulomb interaction U , the fermion operators act in the restricted Hilbert space without double occupancies in the $\{a, b\}$ orbitals which is implemented by the projection operators $\mathcal{P}$. Then: (i) itinerant electrons in yz or zx degenerate orbitals at site j can hop in the allowed $[(\hat{y}, \hat{z}) \text{ or } (\hat{x}, \hat{z})]$ plane to the nearest neighbor (nn) site i , but *only* if there are no other electrons at site i in these orbitals (first and second term), (ii) Hund's exchange $\propto J_H$ aligns spins of itinerant electrons with "core spins" (third terms), and (iii) occupied xy orbitals are responsible for AF SE interactions between nn sites in the $(\hat{x}, \hat{y})$ plane (last term), while the SE interactions due to itinerant electrons are not included.

The above simplifications in the SE part $\propto J$ are motivated by the experimental observation that the structural transitions (which favor the orbital order and occupied xy orbitals) occur in $\text{La}_{1-x}\text{Sr}_x\text{VO}_3$ at lower temperature than the magnetic ones in the entire doping range [3]. Note that the structural transition does not

occur in the metallic phase and then one does not expect any orbital order, similar to the FM metallic manganites [7]. Note that the SE interactions due to the excitations of yz and zx electrons could in principle be included, and would: (i) enhance the AF interactions and induce their alternating orbital (AO) order in $(\hat{x}, \hat{y})$ planes, and (ii) would give FM interactions along the $\hat{z}$ direction, and thus further stabilize the C-AF phase [2].

As in Ref. [9], we treat "core spins" $S = 1/2$ classically and we make a local unitary transformation of the basis $\{a_{i\uparrow}^\dagger|0\rangle, a_{i\downarrow}^\dagger|0\rangle\} \otimes \{b_{i\uparrow}^\dagger|0\rangle, b_{i\downarrow}^\dagger|0\rangle\}$ in the Hilbert space. The transformation makes the quantization axes of the spin of an itinerant electron locally parallel to the one of the "core spin", and thus it diagonalizes Hund's term in the spin variables. Assuming also infinite Hund's exchange $J_H \rightarrow \infty$, we leave out the itinerant electrons with antiparallel spin and what is left is a renormalized (*a priori* nonhomogenous) hopping $t \rightarrow tu_{ij}$ of spinless electrons. The absolute value of the complex parameter u_{ij} is $|u_{ij}| = \cos(\theta_{ij}/2)$, with θ_{ij} being the relative angle between "core spins" at site i and j (the phase of u_{ij} can be chosen arbitrarily at $T = 0$). Furthermore, we will discuss only uniform magnetic structures and follow the so-called uniform hopping approach (UHA) [9]. One obtains then the following *classical* (i.e. Zener-type [6]) DE Hamiltonian:

$$\begin{aligned} \mathcal{H}_{\text{DE}} = & -tu_y \sum_{i,j \parallel \hat{y}} \tilde{a}_i^\dagger \tilde{a}_j - t \sum_{i,j \parallel \hat{z}} \tilde{a}_i^\dagger \tilde{a}_j - tu_x \sum_{i,j \parallel \hat{x}} \tilde{b}_i^\dagger \tilde{b}_j - t \sum_{i,j \parallel \hat{z}} \tilde{b}_i^\dagger \tilde{b}_j \\ & + \frac{J}{2} L^3 (u_x^2 + u_y^2 - 1), \end{aligned} \quad (2)$$

where: $u_x = \cos(\theta_x/2)$, $u_y = \cos(\theta_y/2)$, θ_x (θ_y) is the relative angle between spins in the $\hat{x}$ ($\hat{y}$) direction and L^3 is the number of sites in the crystal. The restricted fermion creation operators are now $\tilde{a}_i^\dagger = a_i^\dagger(1 - b_i^\dagger b_i)$, $\tilde{b}_i^\dagger = b_i^\dagger(1 - a_i^\dagger a_i) - a_i^\dagger(b_i^\dagger)$ creates a spinless electron at site i in $yz(zx)$ orbital. The FM order along the $\hat{z}$ direction is already assumed in Eq. (2) as it minimizes the total energy of the system by maximalizing the possibility of hopping in this direction. Note that all but one (spin 1/2 being classical) of the assumptions necessary to derive Eq. (2) from Eq. (1) seem to be reasonable approximations for cubic vanadates *and* for investigation of the problems addressed in section 1. We leave quantum treatment of the DE for further studies but we believe that the main features of the DE mechanism in cubic vanadates can be captured by this classical Hamiltonian.

3 Numerical results and discussion

The ground state of Eq. (2) was found by introducing Kotliar-Ruckenstein slave boson representation [10]. Following Ref. [7], we adopted this method to the orbital case and made the mean-field approximation (the unrestricted slave boson method, identical with the Gutzwiller approximation and called SBMF). It leads to the effective hopping terms in $\mathcal{H}_{\text{DE}}$:

$$t \tilde{a}_i^\dagger \tilde{a}_j \rightarrow \frac{x}{x_a} t c_i^\dagger c_j, \quad t \tilde{b}_i^\dagger \tilde{b}_j \rightarrow \frac{x}{x_b} t d_i^\dagger d_j, \quad (3)$$




Figure 1: Magnetic phase diagram at $T = 0$ obtained by solving Eq. (2) within SBMF approximation. Under increasing hole doping x , the C -AF order changes via the canted Ca^{**} phase, with spins canted on two bonds, to a homogenous FM phase.

where: $c_i^\dagger$ ($d_i^\dagger$) are pseudofermion creation operators for yz (zx) orbital, and x_a (x_b) are hole numbers in these orbitals with the constraint $x_a + x_b = 1 + x$. Now Eq. (2) is exactly solvable by minimizing the total energy with respect to the u_x and u_y variational parameters [11], which determine the magnetic structure. The resulting magnetic phases are shown in Fig. 1 for different values of SE J/t and hole doping x . We should stress that equal number of holes in both orbitals ($x_a = x_b$) is always concomitant with the obtained magnetic phases, though we did not impose such a condition. This result follows almost entirely from the fermion properties of the system, namely the condition that the Fermi energies of the systems with electrons filling partly yz orbitals and zx orbitals should be the same. Note also that $x_a = x_b$ is a necessary condition for the existence of the AO order. Thus, including real SE interactions from section 1: (i) would not contradict our results in the orbital sector, (ii) would only further support the stability of the C -AF phase in the spin sector, which is already stable for a broad range of parameters.

Let us further analyze these results. One finds that for the broad range of realistic values of parameters, the DE mechanism does not win in $(\hat{x}, \hat{y})$ planes, which instead have AF order and are insulating, while the holes hop only along the FM $\hat{z}$ direction. In this way the C -AF order characterized by the *one-dimensional metallic* behavior is stabilized. Remarkably, such a phase is stable for $0.178 < x < 0.26$ and $J/t = 0.178$, in agreement with the experimental observations [3]. These results are striking since at the first sight one expects that the DE should win here due to the small "core spin" value $S = 1/2$, much smaller than in the manganites ($S = 3/2$). Instead we get results which are qualitatively similar to those obtained by van den Brink and Khomskii in the DE model for electron doped SrMnO_3 [11]. Though, the reason is different — this behavior here follows from very specific features of t_{2g} orbitals: (i) the AF SE interactions are strictly two-dimensional (2D) since for the occupied xy electrons virtual hopping processes (leading to the SE) can occur only in the $(\hat{x}, \hat{y})$ plane — this enables metallic behavior without




Figure 2: Magnetic phase diagrams at $T = 0$, with displayed magnetic order along the $\hat{x}$ axis, as obtained by solving Eq. (2) and assuming: (a) no electrons in yz orbital, and (b) *in addition* no hopping along the $\hat{z}$ direction. The AF order for $x = 0$ [marked in grey (red online)] was found by finite size scaling of the results obtained for large clusters. Note that the AF (FM) order is stable in the entire range of x along the $\hat{y}$ ($\hat{z}$) axis [though the type of order along $\hat{z}$ direction does not matter in (b)].

any loss of the magnetic energy (hopping in the $\hat{z}$ direction does not destroy the magnetic order); (ii) in addition, the itinerant electrons hop between *the other two* t_{2g} orbitals, allowing for the coexistence of the magnetic order and the metallic behavior, and costing no extra magnetic energy.

To understand better this behavior we show in Fig. 2(a) stable magnetic structures in the $\hat{x}$ direction for the systems with 2D planar ($\hat{x}$, $\hat{y}$) AF interactions, and holes being doped *only* into zx orbital, i.e., assuming empty yz orbitals, $\forall_i a_i^\dagger a_i = 0$, in Eq. (2). In addition, we also studied the above case but when the hopping between zx orbitals occurs only along the $\hat{x}$ direction, which resembles holes doped into $d_{3x^2-r^2}$ orbitals in the manganites [stable phases are displayed in Fig. 2(b)]. One finds that for this e_g -type orbital the canted (Ca) structure is stabilized in a broad range of parameters, in contrast with t_{2g} case where AF order in the plane can be conserved. This is one of the reasons why the AF order along the $\hat{z}$ direction can be so easily destabilized by doping in LaMnO_3 , leading to the FM metallic phase [8]. By looking at Fig. 2(a), one can also realize better the role of orbital degeneracy: even if we had a system with planar AF interactions and we doped it with holes into, e.g., zx orbitals, we would not get the AF plane stable for a *realistic* range of parameters! Instead, one AF bond has to be broken. This tendency is suppressed if we have two degenerate orbitals as in Eq. (2).

In summary, we have shown that the existence of the metallic C-AF phase in doped $\text{La}_{1-x}\text{Sr}_x\text{VO}_3$ can be explained by the DE model with extremely simplified SE interactions. The stability of this phase is due to the specific features of the t_{2g} orbitals — namely due to strictly 2D and strictly flavor conserving hopping between these degenerate orbitals [2]. In our view these are very prospective results and, in connection with those of Ref. [5], they suggest that including the real

SE interactions could indeed explain the observed insulating phase and the metal-insulator transition which occurs at finite doping.

Acknowledgments We thank G. Khaliullin for insightful discussions. We would like to acknowledge support by the Polish Ministry of Science and Education under Project No. 1 P03B 068 26.

References

- [1] S. Miyasaka, Y. Okimoto, M. Iwama, and Y. Tokura, Phys. Rev. B **68**, 100406(R) (2003).
- [2] G. Khaliullin, P. Horsch, and A. M. Oleś, Phys. Rev. Lett. **86**, 3879 (2001).
- [3] S. Miyasaka, T. Okuda, and Y. Tokura, Phys. Rev. Lett. **85**, 5388 (2000).
- [4] T. Mizokawa and A. Fujimori, Phys. Rev. B **54**, 5368 (1996).
- [5] S. Ishihara, Phys. Rev. Lett **94**, 156408 (2005).
- [6] C. Zener, Phys. Rev. **82**, 403 (1951).
- [7] L. F. Feiner and A. M. Oleś, Phys. Rev. B **71**, 144422 (2005).
- [8] E. Dagotto, T. Hotta, and A. Moreo, Phys. Rep. **344**, 1 (2001).
- [9] W. Koller, A. Prüll, H. G. Evertz, and W. von der Linden, Phys. Rev. B **66**, 144425 (2002).
- [10] G. Kotliar and A. E. Ruckenstein, Phys. Rev. Lett. **57**, 1362 (1986).
- [11] J. van den Brink and D. I. Khomskii, Phys. Rev. Lett. **82**, 1016 (1999).